

Dr. Kelly McCracken (913) 788-7600 www.amazingsmileskc.com

POST-OPERATIVE INSTRUCTIONS FOR ENDODONTIC PATIENTS

You have received Endodontic treatment in an effort to retain your tooth and avoid extractions. Under most circumstances, about 90% of our treatments are successful in achieving this goal. However, there are some possible complications of which you should be aware.

- 1. Even though the nerve has been removed from the tooth, the act of performing treatment causes the ligament which surrounds the tooth to become inflamed. Most patients report a pain level of "3" on a scale of 1 to 10. This is normal and does not necessarily mean that something is wrong.
- 2. Please take 3 (three) 200mg Ibuprofen/Advil tablets four times a day (not to exceed 2400mg/day) for three days to control this discomfort. If you are unable to take Advil, take 2 (two) Extra Strength Tylenol 500mg tablets four times per day (not to exceed 4000mg/day) for three days.
- 3. If this does not control your discomfort, fill the prescription (if given) as directed. If you did not receive a pain control prescription and are experiencing pain that is not comforted with Advil/Tylenol, please call our office immediately. You may supplement the prescription with the above mentioned Advil/Tylenol advice for added pain control. Be aware that your pain medication may already contain acetaminophen (Tylenol) so do not exceed the above mentioned recommendation. If you have a liver condition, please let the office/pharmacist know.
- 4. It is normal for a tooth that has had endodontic treatment to be sore to biting pressure, so please avoid chewing on the tooth for at least one week.
- 5. Approximately 2% of root canal treatments will "flare up." This means that the symptoms become worse instead of better. If your pain level is increasing rather than decreasing and/or swelling develops, please contact our office. If you are calling after-hours, the doctor's cell phone is stated on the office's answering message.
- 6. You will need to schedule your final restoration within 2-4 weeks of completion of root canal treatment. While your pain may have subsided, the tooth is in a temporary state and the remaining tooth structure is fragile and unsupported. Without a final restoration your root canal treatment may be compromised and may need to be redone in the future.

If you have any questions, don't hesitate to contact us! We are here to help you and wish the best for your dental health.